

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:ssptacrs1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 DEC 01 ChemPort single article sales feature unavailable
NEWS 3 APR 03 CAS coverage of exemplified prophetic substances enhanced
NEWS 4 APR 07 STN is raising the limits on saved answers
NEWS 5 APR 24 CA/CAPLUS now has more comprehensive patent assignee information
NEWS 6 APR 26 USPATFULL and USPAT2 enhanced with patent assignment/reassignment information
NEWS 7 APR 28 CAS patent authority coverage expanded
NEWS 8 APR 28 ENCOMPLIT/ENCOMPLIT2 search fields enhanced
NEWS 9 APR 28 Limits doubled for structure searching in CAS REGISTRY
NEWS 10 MAY 08 STN Express, Version 8.4, now available
NEWS 11 MAY 11 STN on the Web enhanced
NEWS 12 MAY 11 BEILSTEIN substance information now available on STN Easy
NEWS 13 MAY 14 DGENE, PCTGEN and USGENE enhanced with increased limits for exact sequence match searches and introduction of free HIT display format
NEWS 14 MAY 15 INPADOCDB and INPAFAMDB enhanced with Chinese legal status data
NEWS 15 MAY 28 CAS databases on STN enhanced with NANO super role in records back to 1992
NEWS 16 JUN 01 CAS REGISTRY Source of Registration (SR) searching enhanced on STN
NEWS 17 JUN 26 NUTRACEUT and PHARMAML no longer updated
NEWS 18 JUN 29 IMSCOPROFILE now reloaded monthly
NEWS 19 JUN 29 EFFULL adds Simultaneous Left and Right Truncation (SLART) to AB, MCLM, and TI fields
NEWS 20 JUL 09 PATDPAFULL adds Simultaneous Left and Right Truncation (SLART) to AB, CLM, MCLM, and TI fields
NEWS 21 JUL 14 USGENE enhances coverage of patent sequence location (PSL) data
NEWS 22 JUL 27 CA/CAPLUS enhanced with new citing references
NEWS 23 JUL 16 GBFULL adds patent backfile data to 1855
NEWS 24 JUL 21 USGENE adds bibliographic and sequence information

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 07:40:47 ON 28 JUL 2009

=> file registry

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.22

0.22

FILE 'REGISTRY' ENTERED AT 07:41:08 ON 28 JUL 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 26 JUL 2009 HIGHEST RN 1169218-44-3

DICTIONARY FILE UPDATES: 26 JUL 2009 HIGHEST RN 1169218-44-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Program Files\STNEXP\Queries\10588454.str



chain nodes :

7 8 9 10 11 12 13 14 15 16 17 18 19

ring nodes :

1 2 3 4 5 6

chain bonds :

2-7 5-11 7-8 8-9 8-10 11-12 11-13 13-14 13-16 14-15 15-17 15-18 15-19

ring bonds :
 1-2 1-6 2-3 3-4 4-5 5-6
 exact/norm bonds :
 13-14 13-16 14-15 15-17 15-18 15-19
 exact bonds :
 2-7 5-11 7-8 8-9 8-10 11-12 11-13
 normalized bonds :
 1-2 1-6 2-3 3-4 4-5 5-6


Match level :
 1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS 18:CLASS
 19:CLASS

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1 fam ful
 FULL SEARCH INITIATED 07:41:26 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 59 TO ITERATE

100.0% PROCESSED 59 ITERATIONS 5 ANSWERS
 SEARCH TIME: 00.00.01

L2 5 SEA FAM FUL L1

=> file caplus
 COST IN U.S. DOLLARS
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
73.33	73.55

FILE 'CAPLUS' ENTERED AT 07:41:46 ON 28 JUL 2009
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available

for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 28 Jul 2009 VOL 151 ISS 5
 FILE LAST UPDATED: 27 Jul 2009 (20090727/ED)
 REVISED CLASS FIELDS (/NCL) LAST RELOADED: Jun 2009
 USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Jun 2009

CAplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

The ALL, BIB, MAX, and STD display formats in the CA/CAplus family of databases have been updated to include new citing references information. This enhancement may impact record import into database management software. For additional information, refer to NEWS 22.

```
=> s 12
L3          23 L2

=> s 13 and spin?
          746103 SPIN?
L4          3 L3 AND SPIN?

=> d 14 ibib abs 1-3
```

```
L4 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2008:412188 CAPLUS
DOCUMENT NUMBER: 148:394429
TITLE: CXC chemokine-mediated signaling targeting for
treatment of a myelin disorder
INVENTOR(S): Miller, Robert H.; Padovani-Claudio, Dolly A.
PATENT ASSIGNEE(S): Case Western Reserve University, USA
SOURCE: PCT Int. Appl., 85pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:
```

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008039876	A1	20080403	WO 2007-US79602	20070926
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,				
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI,				
GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG,				
KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,				
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,				
PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,				
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,
GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM

CA 2664359 A1 20080403 CA 2007-2664359 20070926
US 20090041753 A1 20090212 US 2007-904634 20070926
EP 2066335 A1 20090610 EP 2007-843271 20070926

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
IS, IT, LI, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR,
AL, BA, HR, MK, RS

PRIORITY APPLN. INFO.: US 2006-847656P P 20060926
WO 2007-US79602 W 20070926

AB The invention discloses compns. and methods for targeting CXCR2
chemokine-mediated signaling for treatment of a myelin disorder. The
methodol. of the invention can be used to ameliorate neuropathies.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2009 ACS ON SIN

ACCESSION NUMBER: 2007:976827 CAPLUS

DOCUMENT NUMBER: 147:314799

TITLE: Reparixin, an inhibitor of CXCR2 function, attenuates
inflammatory responses and promotes recovery of
function after traumatic lesion to the spinal
cord

AUTHOR(S): Gorio, Alfredo; Madaschi, Laura; Zadra, Georgia;
Marfia, Giovanni; Cavallieri, Barbara; Bertini,
Riccardo; Di Giulio, Anna Maria

CORPORATE SOURCE: Pharmacological Laboratories, Department of Medicine,
Surgery and Dentistry, Faculty of Medicine, University
of Milan, Milan, Italy

SOURCE: Journal of Pharmacology and Experimental Therapeutics
(2007), 322(3), 973-981
CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: American Society for Pharmacology and Experimental
Therapeutics

DOCUMENT TYPE: Journal

LANGUAGE: English

AB It has been shown that the blockade of CXCR1 and CXCR2 receptors prevents
ischemia/reperfusion damage in several types of vascular beds. Reparixin
is a recently described inhibitor of human CXCR1/R2 and rat CXCR2 receptor
activation. We applied reparixin in rats following traumatic
spinal cord injury and determined therapeutic temporal and dosages
windows. Treatment with reparixin significantly counteracts secondary
degeneration by reducing oligodendrocyte apoptosis, migration to the
injury site of neutrophils and ED-1-pos. cells. The observed preservation of
the white matter might also be secondary to the enhanced proliferation of
NG2-pos. cells. The expression of macrophage-inflammatory protein-2,
tumor necrosis factor- α , interleukin (IL)-6, and IL-1 β was also
counteracted, and the proliferation of glial fibrillary acidic
protein-pos. cells was markedly reduced. These effects resulted in a
smaller post-traumatic cavity and in a significantly improved recovery of
hind limb function. The best beneficial outcome of reparixin treatment
required 7-day administration either by i.p. route (15 mg/kg) or s.c.
infusion via osmotic pumps (10 mg/kg), reaching a steady blood level of 8
 μ g/mL. Methylprednisolone was used as a reference drug; such treatment
reduced cytokine production but failed to affect the rate of hind limb
recovery.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD
(4 CITINGS)

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:704377 CAPLUS

DOCUMENT NUMBER: 145:369213

TITLE: Species differences in the pharmacokinetics and metabolism of Reparixin in rat and dog

AUTHOR(S): Midgley, I.; Fitzpatrick, K.; Wright, S. J.; John, B. A.; Peard, A. J.; Major, R. M.; Holding, J. D.; McBurney, A.; Anacardio, R.; Novellini, R.; Ferrari, M. P.

CORPORATE SOURCE: Department of Drug Metabolism, Huntingdon Life Sciences Ltd, Huntingdon, UK

SOURCE: Xenobiotica (2006), 36(5), 419-440
CODEN: XENOBH; ISSN: 0049-8254

PUBLISHER: Taylor & Francis Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The pharmacokinetics and metabolism of Reparixin (formerly Repertaxin), a potent and specific inhibitor of the chemokine CXCL8, were investigated in rats and dogs after i.v. administration of [14C]Reparixin L-lysine salt. Protein binding of Reparixin was investigated in vitro in rat, dog, rabbit, cynomolgus monkey, and human plasma. Plasma protein binding of Reparixin was >99% in the laboratory animals and humans up to 50 µg mL⁻¹, but lower at higher concns. Although radioactivity was rapidly distributed into rat tissues, V_{ss} was low (.apprx.0.15 L kg⁻¹) in both rat and dog. Nevertheless, Reparixin was more rapidly eliminated in rats (t_{1/2} .apprx.0.5 h) than in dogs (t_{1/2} .apprx.10 h). Systemic exposure in dog was due primarily to parent drug, but metabolites played a more prominent role in rat. Oxidation of the iso-Bu side-chain was the major metabolic pathway in rat, whereas hydrolysis of the amide bond predominated in dog. Urinary excretion, which accounted for 80-82% of the radioactive dose, was the major route of elimination in both species, and biotransformation of Reparixin was complete before excretion.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d 13 1-23 ibib abs

L3 ANSWER 1 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2009:779262 CAPLUS

TITLE: Development and validation of an LC-MS/MS method for determination of methanesulfonamide in human urine

AUTHOR(S): Anacardio, Roberto; Mullins, Frank G. P.; Hannam, Sally; Sheikh, Muhammed S.; O'Shea, Karen; Aramini, Andrea; D'Anniballe, Gaetano; D'Anteo, Loredana; Ferrari, Mauro P.; Allegretti, Marcello

CORPORATE SOURCE: Research Department, Dompe pha.r.ma s.p.a., L'Aquila, Italy

SOURCE: Journal of Chromatography, B: Analytical Technologies in the Biomedical and Life Sciences (2009), 877(22), 2087-2092

CODEN: JCBAAI; ISSN: 1570-0232

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A sensitive and selective liquid chromatog. method coupled with tandem mass spectrometry (LC-MS/MS) was developed and validated for the quantification of methanesulfonamide (MSA) in human urine. MSA is a potential in vivo

metabolite of reparixin, a specific inhibitor of the CXCL8 biol. activity. In this study, a simple derivatization procedure with a new reagent, N-(4-methanesulfonyl-benzoyl)-imidazole, was set up to enable MSA and the internal standard (I.S.), ethanesulfonamide (ESA), to be analyzed by LC-MS/MS. After derivatization, samples were evaporated and reconstituted in 30% acetonitrile, aqueous MSA and I.S. derivs. were separated by reversed phased

HPLC

(high performance liquid chromatog.) on a Luna 5 μ C18 column and quantitated by MS/MS using electrospray ionization (ESI) and multiple reaction monitoring (MRM) in the neg. ion mode. The most intense [M-H]⁻MRM transition of derivatized MSA at m/z 276.2 \rightarrow 197.2 was used for quantitation and the transition at m/z 290.2 \rightarrow 211.2 was used to monitor derivatized ESA. The method was linear over the concentration range

from

1 to 100 μ g/mL, with a lower limit of quantitation of 1 μ g/mL. The intra- and inter-day precisions were less than 5.5% and 10.1%, resp., and the accuracies were between -4.0% and +11.3%. The method was successfully applied to quantify levels of MSA in human urine after i.v. administration of reparixin to healthy volunteers.

L3 ANSWER 2 OF 23 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2008:1475435 CAPLUS

DOCUMENT NUMBER: 150:75537

TITLE: Novel Role of CXCR2 in Regulation of γ -Secretase Activity

AUTHOR(S): Bakshi, Pancham; Margenthaler, Elaina; Laporte, Vincent; Crawford, Fiona; Mullan, Michael

CORPORATE SOURCE: Roskamp Institute, Sarasota, FL, 34203, USA

SOURCE: ACS Chemical Biology (2008), 3(12), 777-789

CODEN: ACBCCT; ISSN: 1554-8929

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Alzheimer's disease (AD) is a progressive chronic disorder that leads to cognitive decline. Several studies have associated up-regulation of some of the chemokines and/or their receptors with altered APP processing leading to increased production of β -amyloid protein (β) and AD pathol. changes. However, there is no direct evidence to date to determine whether the altered processing of APP results in up-regulation of these receptors or whether the up-regulation of the chemokine receptors causes modulated processing of APP. In the current study, we demonstrate that treatment of the chemokine receptor CXCR2 with agonists leads to enhancement of β production and treatment with antagonists or immunodepletion of CXCR2's endogenous agonists leads to β inhibition. Further, we found that the inhibitory effect of the antagonist of CXCR2 on β 40 and β 42 is mediated via γ -secretase, specifically through reduction in expression of presenilin (PS), one of the γ -secretase components. Also, in vivo chronic treatment with a CXCR2 antagonist blocked β 40 and β 42 production. Using small interfering RNAs for CXCR2, we further showed that knockdown of CXCR2 in vitro accumulates γ -secretase substrates C99 and C83 with reduced production of both β 40 and β 42. Taken together, these findings strongly suggest for the first time that up-regulation of the CXCR2 receptor can be the driving force in increased production of β . Our findings unravel new mechanisms involving the CXCR2 receptor in the pathogenesis of AD and pose it as a potential target for developing novel therapeutics for intervention in this disease. Also, we propose here a new chemical series of interest that can serve as a prototype for drug development.

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 3 OF 23 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2008:1167893 CAPLUS
 DOCUMENT NUMBER: 149:439943
 TITLE: Therapeutic inhibition of CXCR2 by Reparixin attenuates acute lung injury in mice
 AUTHOR(S): Zarbock, A.; Allegretti, M.; Ley, K.
 CORPORATE SOURCE: Division of Inflammation Biology, La Jolla Institute for Allergy and Immunology, La Jolla, CA, USA
 SOURCE: British Journal of Pharmacology (2008), 155(3), 357-364
 PUBLISHER: CODEN: BJPCBM; ISSN: 0007-1188
 DOCUMENT TYPE: Nature Publishing Group
 LANGUAGE: Journal
 AB English

AB Acute lung injury (ALI) remains a major challenge in critical care medicine. Both neutrophils and chemokines have been proposed as key components in the development of ALI. The main chemokine receptor on neutrophils is CXCR2, which regulates neutrophil recruitment and vascular permeability, but no small mol. CXCR2 inhibitor has been demonstrated to be effective in ALI or animal models of ALI. To investigate the functional relevance of the CXCR2 inhibitor reparixin in vivo, we determined its effects in two models of ALI, induced by either lipopolysaccharide (LPS) inhalation or acid instillation. In two ALI models in mice, we measured vascular permeability by Evans blue and evaluated neutrophil recruitment into the lung vasculature, interstitium and airspace by flow cytometry. Pharmacol. inhibition of CXCR2 by reparixin reduced CXCL1-induced leukocyte arrest in the microcirculation of the cremaster muscle, but did not influence arrest in response to leukotriene B4 (LTB4) demonstrating specificity. Reparixin (15 µg g-1) reduced neutrophil recruitment in the lung by approx. 50% in a model of LPS-induced ALI. A higher dose did not provide addnl. reduction of neutrophil recruitment. This dose also reduced accumulation of neutrophils in the interstitial compartment and vascular permeability in LPS-induced ALI. Furthermore, both prophylactic and therapeutic application of reparixin improved gas exchange, and reduced neutrophil recruitment and vascular permeability in a clin. relevant model of acid-induced ALI. Reparixin, a non-competitive allosteric CXCR2 inhibitor attenuates ALI by reducing neutrophil recruitment and vascular permeability.

REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:589691 CAPLUS
 DOCUMENT NUMBER: 148:554109
 TITLE: Method and use of nonionic polymers for increasing efficacy of anti-adhesive compositions in controlling inflammation and pain
 INVENTOR(S): Chamness, Kathy L.
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 15pp.
 DOCUMENT TYPE: CODEN: USXXCO
 LANGUAGE: Patent
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION: English

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
US 20080112921	A1	20080515	US 2006-598397	20061114
WO 2008063943	A2	20080529	WO 2007-US84387	20071112
WO 2008063943	A3	20090507		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI,

GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG,
 KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,
 PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,
 GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
 BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA

PRIORITY APPLN. INFO.: US 2006-598397 A 20061114

AB The invention discloses a method and kits for increasing the efficiency of anti-adhesive compns. by parenterally administering a composition comprising an effective amount of at least one pharmaceutically acceptable anti-adhesive nonionic polymer to a site of injury, controlling inflammation at the site of injury, and reducing pain. The nonionic polymers are used with magnesium salts.

L3 ANSWER 5 OF 23 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2008:412188 CAPLUS

DOCUMENT NUMBER: 148:394429

TITLE: CXC chemokine-mediated signaling targeting for treatment of a myelin disorder

INVENTOR(S): Miller, Robert H.; Padovani-Claudio, Dolly A.

PATENT ASSIGNEE(S): Case Western Reserve University, USA

SOURCE: PCT Int. Appl., 85pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008039876	A1	20080403	WO 2007-US79602	20070926
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
CA 2664359	A1	20080403	CA 2007-2664359	20070926
US 20090041753	A1	20090212	US 2007-904634	20070926
EP 2066335	A1	20090610	EP 2007-843271	20070926
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS			

PRIORITY APPLN. INFO.: US 2006-847656P P 20060926

WO 2007-US79602 W 20070926

AB The invention discloses compns. and methods for targeting CXC chemokine-mediated signaling for treatment of a myelin disorder. The methodol. of the invention can be used to ameliorate neuropathies.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 6 OF 23 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2007:1075862 CAPLUS
 DOCUMENT NUMBER: 147:541555
 TITLE: A new and efficient method for the facile synthesis of N-acyl sulfonamides under Lewis acid catalysis
 AUTHOR(S): Reddy, Chada Raji; Mahipal, Budugam; Yaragorla, Srinivasa Rao
 CORPORATE SOURCE: Organic Division-I, Indian Institute of Chemical Technology, Hyderabad, 500 007, India
 SOURCE: Tetrahedron Letters (2007), 48(42), 7528-7532
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 147:541555

AB The N-acylation of sulfonamides with carboxylic acid anhydrides in the presence of Lewis acids is described. Several Lewis acids such as BF₃·Et₂O, ZnCl₂, MoCl₅, TiCl₄, B(C₆F₅)₃, Sc(OTf)₃ and I₂ were found to catalyze the reaction efficiently to furnish the N-acylated products in good yields under solvent-free conditions. The reactions of various sulfonamides were studied with different carboxylic acid anhydrides including the less reactive benzoic and pivalic anhydrides, in the presence of 3 mol% ZnCl₂ as the catalyst. Carboxylic acids were also successfully used as acylating agents via the mixed anhydride method.
 OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)
 REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 7 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 2007:993886 CAPLUS
 DOCUMENT NUMBER: 147:292200
 TITLE: Methods and compositions for treating and preventing tumors
 INVENTOR(S): Bonni, Azad M.; De la Iglesia, Nuria; Konopka, Genevieve
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 21pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070208074	A1	20070906	US 2007-657965	20070124
PRIORITY APPLN. INFO.:			US 2006-762033P	P 20060124
AB The present invention provides methods for reducing the growth or invasiveness of tumors.				

L3 ANSWER 8 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 2007:976827 CAPLUS
 DOCUMENT NUMBER: 147:314799
 TITLE: Reparixin, an inhibitor of CXCR2 function, attenuates inflammatory responses and promotes recovery of function after traumatic lesion to the spinal cord
 AUTHOR(S): Gorio, Alfredo; Madaschi, Laura; Zadra, Giorgia; Marfia, Giovanni; Cavalieri, Barbara; Bertini, Riccardo; Di Giulio, Anna Maria
 CORPORATE SOURCE: Pharmacological Laboratories, Department of Medicine, Surgery and Dentistry, Faculty of Medicine, University of Milan, Milan, Italy

SOURCE: Journal of Pharmacology and Experimental Therapeutics
(2007), 322(3), 973-981

CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: American Society for Pharmacology and Experimental
Therapeutics

DOCUMENT TYPE: Journal

LANGUAGE: English

AB It has been shown that the blockade of CXCR1 and CXCR2 receptors prevents ischemia/reperfusion damage in several types of vascular beds. Reparixin is a recently described inhibitor of human CXCR1/R2 and rat CXCR2 receptor activation. We applied reparixin in rats following traumatic spinal cord injury and determined therapeutic temporal and dosages windows. Treatment with reparixin significantly counteracts secondary degeneration by reducing oligodendrocyte apoptosis, migration to the injury site of neutrophils and ED-1-pos. cells. The observed preservation of the white matter might also be secondary to the enhanced proliferation of NG2-pos. cells. The expression of macrophage-inflammatory protein-2, tumor necrosis factor- α , interleukin (IL)-6, and IL-1 β was also counteracted, and the proliferation of glial fibrillary acidic protein-pos. cells was markedly reduced. These effects resulted in a smaller post-traumatic cavity and in a significantly improved recovery of hind limb function. The best beneficial outcome of reparixin treatment required 7-day administration either by i.p. route (15 mg/kg) or s.c. infusion via osmotic pumps (10 mg/kg), reaching a steady blood level of 8 μ g/mL. Methylprednisolone was used as a reference drug; such treatment reduced cytokine production but failed to affect the rate of hind limb recovery.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD
(4 CITINGS)

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 9 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:807542 CAPLUS

DOCUMENT NUMBER: 147:314717

TITLE: The interleukin-8 (IL-8/CXCL8) receptor inhibitor
reparixin improves neurological deficits and reduces
long-term inflammation in permanent and transient
cerebral ischemia in rats

AUTHOR(S): Villa, Pia; Triulzi, Sara; Cavalieri, Barbara; Di
Bitondo, Rosa; Bertini, Riccardo; Barbera, Sara;
Bigini, Paolo; Mennini, Tiziana; Gelosa, Paolo;
Tremoli, Elena; Sironi, Luigi; Ghezzi, Pietro

CORPORATE SOURCE: Mario Negri Institute, Milan, 20157, Italy
SOURCE: Molecular Medicine (Manhasset, NY, United States)
(2007), 13(3-4), 125-133
CODEN: MOMEF3; ISSN: 1076-1551

PUBLISHER: Feinstein Institute for Medical Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Leukocyte infiltration is viewed as a pharmacol. target in cerebral ischemia. We previously reported that reparixin, a CXCL8 receptor blocker that inhibits neutrophil infiltration, and related mols. can reduce infarct size in a rat model of transient middle cerebral artery occlusion (MCAO). The study aims were to compare the effects of reparixin in transient and permanent MCAO using varied treatment schedules and therapeutic windows to evaluate effects on long-term neurol. deficits and late inflammatory response. Reparixin, administered for 1 to 3 days, 3.5 to 6 h after MCAO, ameliorates neurol. function recovery and inhibits long-term inflammation. The infarct size reduction at 24 h, evaluated by TTC staining, is more pronounced in transient MCAO. MRI anal. identified a decrease in the progression of infarct size by reparixin that was more

evident at 48 h in permanent MCAO, and was associated with a significantly improved recovery from long-term neurol. deficits.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 10 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:451972 CAPLUS

DOCUMENT NUMBER: 147:109107

TITLE: Reparixin, a specific interleukin-8 inhibitor, has no effects on inflammation during endotoxemia

AUTHOR(S): Leitner, J. M.; Mayr, F. B.; Firbas, C.; Spiel, A. O.; Steinlechner, B.; Novellini, R.; Jilma, B.

CORPORATE SOURCE: Department of Clinical Pharmacology, Division of Immunoheumatology, Medical University of Vienna, Austria

SOURCE: International Journal of Immunopathology and Pharmacology (2007), 20(1), 25-36

CODEN: IJIFE4; ISSN: 0394-6320

PUBLISHER: Biolife s.a.s.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Reparixin antagonizes interleukin-8 (IL-8) on the level of signal transduction in vitro. We hypothesized that IL-8 mediates some of the reactions occurring during acute inflammation and specifically that IL-8 may be a mediator of endotoxin induced neutrophilia. We therefore tested the effects of reparixin on humoral and cellular parameters in LPS-induced acute systemic inflammation. The study is a randomized (3:2 active:placebo), double-blind, placebo-controlled parallel group trial. Twenty healthy male volunteers randomly received either reparixin (12) or placebo (8) i.v. One hour after the start of reparixin/placebo infusion a bolus of 2 ng/kg endotoxin was infused over 1-2 min. Blood samples were obtained over 24 h. Reparixin, being metabolized to ibuprofen, suppressed serum thromboxane B2 levels by 78% compared to baseline and control at 8 h. LPS-induced neutrophilia was not significantly affected by reparixin in human volunteers. Consistently, reparixin did not alter the lymphocyte or monocyte counts and had no effect on LPS-induced systemic inflammation as measured by tumor necrosis factor alpha (TNF-α) or interleukin-6 (IL-6) release. Regulation of IL-8 receptors CXCR1 and 2 and the degranulation marker CD11b showed the expected kinetics. Reparixin had no effect on thrombin formation as measured by prothrombin fragment (F1+2). In conclusion, our study showed that reparixin was safe but had no impact on endotoxin induced inflammation. In contrast to previous studies with its metabolite ibuprofen, reparixin does not enhance inflammation in this model.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 11 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:704377 CAPLUS

DOCUMENT NUMBER: 145:369213

TITLE: Species differences in the pharmacokinetics and metabolism of Reparixin in rat and dog

AUTHOR(S): Midgley, I.; Fitzpatrick, K.; Wright, S. J.; John, B. A.; Peard, A. J.; Major, R. M.; Holding, J. D.; McBurney, A.; Anacardio, R.; Novellini, R.; Ferrari, M. P.

CORPORATE SOURCE: Department of Drug Metabolism, Huntingdon Life Sciences Ltd, Huntingdon, UK

SOURCE: Xenobiotica (2006), 36(5), 419-440
 CODEN: XENOBH; ISSN: 0049-8254
 PUBLISHER: Taylor & Francis Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The pharmacokinetics and metabolism of Reparixin (formerly Repertaxin), a potent and specific inhibitor of the chemokine CXCL8, were investigated in rats and dogs after i.v. administration of [¹⁴C]Reparixin L-lysine salt. Protein binding of Reparixin was investigated in vitro in rat, dog, rabbit, cynomolgus monkey, and human plasma. Plasma protein binding of Reparixin was >99% in the laboratory animals and humans up to 50 µg mL⁻¹, but lower at higher concns. Although radioactivity was rapidly distributed into rat tissues, V_{ss} was low (.apprx.0.15 L kg⁻¹) in both rat and dog. Nevertheless, Reparixin was more rapidly eliminated in rats (t_{1/2} .apprx.0.5 h) than in dogs (t_{1/2} .apprx.10 h). Systemic exposure in dog was due primarily to parent drug, but metabolites played a more prominent role in rat. Oxidation of the iso-Bu side-chain was the major metabolic pathway in rat, whereas hydrolysis of the amide bond predominated in dog. Urinary excretion, which accounted for 80-82% of the radioactive dose, was the major route of elimination in both species, and biotransformation of Reparixin was complete before excretion.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 12 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:608541 CAPLUS

DOCUMENT NUMBER: 145:62689

TITLE: Preparation of 2-arylpropionamides for the inhibition of the chemotactic activation induced by C5a

INVENTOR(S): Allegretti, Marcello; Bertini, Riccardo; Beccari, Andrea; Moriconi, Alessio; Aramini, Andrea; Bizzarri, Cinzia; Colotta, Francesco

PATENT ASSIGNEE(S): Dompe' S.p.A., Italy

SOURCE: PCI Int. Appl., 51 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English



FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006063999	A1	20060622	WO 2005-EP56742	20051213
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
AU 2005315591	A1	20060622	AU 2005-315591	20051213
CA 2589495	A1	20060622	CA 2005-2589495	20051213
EP 1856031	A1	20071121	EP 2005-817430	20051213
EP 1856031	B1	20090225		
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				

IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL,
 BA, HR, MK, YU

JP 2008524157	T	20080710	JP 2007-546040	20051213
AT 423760	T	20090315	AT 2005-817430	20051213
ES 2322487	T3	20090622	ES 2005-817430	20051213
MX 2007007133	A	20070808	MX 2007-7133	20070614
US 20080312293	A1	20081218	US 2007-721971	20070615
KR 2007112365	A	20071123	KR 2007-715497	20070706
NO 2007003622	A	20070917	NO 2007-3622	20070713
CN 101184726	A	20080521	CN 2005-80048026	20070810
PRIORITY APPLN. INFO.:			EP 2004-29684	A 20041215
OTHER SOURCE(S):		CASREACT 145:62689; MARPAT 145:62689	WO 2005-EP56742	W 20051213
GI				



AB Title compds. I [Ar = Ph substituted in the meta position by a group selected from alkanoyl, cycloalkanoyl, heteroarylcarbonyl, etc.; R = H, OH, alkyl, etc.] were prepared For example, chlorination of (R)-2-(3-isobutylphenyl)propionic acid, e.g., prepared from 2-[(3-carboxy)phenyl]propionitrile in 3 steps, using thionyl chloride followed by treatment with ammonia afforded compound II. In C5a induced PMNs chemotaxis inhibition assays, compound II exhibited the activity of 50 ± 7% at 10⁻⁷ M. Compds. I are claimed useful for the treatment of sepsis, psoriasis, etc.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 13 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:1301378 CAPLUS

DOCUMENT NUMBER: 144:324102

TITLE: Neutrophil recruitment in the reperfused-injured rat liver was effectively attenuated by repertaxin, a novel allosteric non-competitive inhibitor of CXCL8 receptors: A therapeutic approach for the treatment of post-ischemic hepatic syndromes

AUTHOR(S): Cavalieri, B.; Mosca, M.; Ramadori, P.; Perrelli, M.-G.; De Simone, L.; Colotta, F.; Bertini, R.; Poli, G.; Cutrin, J. C.

CORPORATE SOURCE: Laboratory of Experimental Liver Pathology, Department of Clinical and Biological Sciences, University of Turin, L'Aquila, Italy

SOURCE: International Journal of Immunopathology and Pharmacology (2005), 18(3), 475-486
CODEN: IJIFE4; ISSN: 0394-6320

PUBLISHER: Biolife s.a.s.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Hepatic reperfusion injury represents a crucial problem in several clin. situations including liver transplantation, extensive hepatectomy and hypovolemic shock with resuscitation. Repertaxin is a new non-competitive allosteric blocker of interleukin-8 (CXCL8) receptors, which by locking CXCR1/R2 in an inactive conformation, prevents receptor signaling and polymorphonuclear leukocyte (PMN) chemotaxis. The present study shows that repertaxin dramatically prevents rat post-ischemic hepatocellular necrosis (80% of inhibition) and PMN infiltration (96% of inhibition) at a clin.-relevant time (24 h) of reperfusion. Treatment with repertaxin by continuous infusion is demonstrated to be the optimal route of administration of the compound especially in view of its clin. therapeutic use. Because repertaxin has proven to be safe and well tolerated in different animal studies and in phase I studies in human volunteers, it is in fact a candidate novel therapeutic agent for the prevention and treatment of hepatic post-ischemic injury.

OS.CITING REF COUNT: 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 14 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:460353 CAPLUS

DOCUMENT NUMBER: 143:145782

TITLE: 2-Arylpropionic CXC Chemokine Receptor 1 (CXCR1) Ligands as Novel Noncompetitive CXCL8 Inhibitors

AUTHOR(S): Allegretti, Marcello; Bertini, Riccardo; Cesta, Maria Candida; Bizzarri, Cinzia; Di Bitondo, Rosa; Di Cioccio, Vito; Galliera, Emanuela; Berdini, Valerio; Topai, Alessandra; Zampella, Giuseppe; Russo, Vincenzo; Di Bello, Nicoletta; Nano, Giuseppe; Nicolini, Luca; Locati, Massimo; Fantucci, Piercarlo; Florio, Saverio; Colotta, Francesco

CORPORATE SOURCE: Dompe Research and Development, Dompe S.p.A., L'Aquila, 67100, Italy

SOURCE: Journal of Medicinal Chemistry (2005), 48(13), 4312-4331
CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:145782

AB The CXC chemokine CXCL8/IL-8 plays a major role in the activation and recruitment of polymorphonuclear (PMN) cells at inflammatory sites. CXCL8 activates PMNs by binding the seven-transmembrane (7-TM) G-protein-coupled receptors CXC chemokine receptor 1 (CXCR1) and CXC chemokine receptor 2 (CXCR2). (R)-Ketoprofen (1) was previously reported to be a potent and specific noncompetitive inhibitor of CXCL8-induced human PMNs chemotaxis. The authors report here mol. modeling studies showing a putative interaction site of 1 in the TM region of CXCR1. The binding model was confirmed by alanine scanning mutagenesis and photoaffinity labeling expts. The mol. model driven medicinal chemical optimization of 1 led to a new class of potent and specific inhibitors of CXCL8 biol. activity. Among these, repertaxin was selected as a clin. candidate drug for

prevention of postischemia reperfusion injury.
OS.CITING REF COUNT: 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS
RECORD (23 CITINGS)
REFERENCE COUNT: 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 15 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2005:437986 CAPLUS
DOCUMENT NUMBER: 143:53210
TITLE: Inhibition of the chemokine receptor CXCR2 prevents
kidney graft function deterioration due to
ischemia/reperfusion
AUTHOR(S): Cugini, Daniela; Azzollini, Nadia; Gagliardini, Elena;
Cassis, Paola; Bertini, Riccardo; Colotta, Francesco;
Noris, Marina; Remuzzi, Giuseppe; Benigni, Ariela
CORPORATE SOURCE: Transplant Research Center "Chiara Cucchi de
Alessandri e Gilberto Crespi" Mario Negri Institute
for Pharmacological Research, Bergamo, Italy
SOURCE: Kidney International (2005), 67(5), 1753-1761
CODEN: KDYIA5; ISSN: 0085-2538
PUBLISHER: Blackwell Publishing, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Background: Ischemia/reperfusion (I/R) injury after organ transplantation
is a major cause of delayed graft function. Following I/R, locally
produced CXC chemokines attract and activate granulocytes, which in turn
promote graft damage. Methods: We examined the involvement of granulocyte
recruitment via the CXCR2 pathway in a rat model of 4 h cold ischemia
followed by kidney transplantation. Serum creatinine and intragraft
granulocyte infiltration were monitored in the early phase posttransplant.
A CXCR2 inhibitor, repertaxin, was given to recipients before
transplantation (at -24 h or -8 h or -2 h), immediately before reperfusion
and 2 h later. Results: An increase of granulocyte chemoattractant
CINC-1/interleukin-8 (IL-8) mRNA expression after I/R both in syngeneic
and allogeneic transplantation was associated with a marked infiltration of
granulocytes in renal tissue. In syngeneic transplantation, Lewis rats
given 15 mg/kg repertaxin 24 h before surgery had granulocyte graft
infiltration and serum creatinine levels significantly reduced in respect
to vehicle-treated animals. Intermediate effects were observed with 5 mg/kg,
whereas the dose of 30 mg/kg had toxic effects. We found that reducing
the pretreatment time to 8 h before surgery was still effective.
Prevention of granulocyte infiltration and serum creatinine increase was
also obtained in allogeneic transplantation, when Brown Norway recipients
of Lewis kidneys were given 15 mg/kg repertaxin starting 8 h before
surgery. Conclusion: Repertaxin treatment of the recipient animal was
effective in preventing granulocyte infiltration and renal function
impairment both in syngeneic and in allogeneic settings. The possibility
to modulate I/R injury in this rat model opens new perspectives for
preventing posttransplant delayed graft function in humans.

OS.CITING REF COUNT: 33 THERE ARE 33 CAPLUS RECORDS THAT CITE THIS
RECORD (33 CITINGS)
REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 16 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2005:319144 CAPLUS
DOCUMENT NUMBER: 142:475974
TITLE: Neuroprotection with the CXCL8 inhibitor repertaxin in
transient brain ischemia
AUTHOR(S): Garau, Angela; Bertini, Riccardo; Colotta, Francesco;
Casilli, Federica; Bigini, Paolo; Cagnotto, Alfredo;
Mennini, Tiziana; Ghezzi, Pietro; Villa, Pia

CORPORATE SOURCE: "Mario Negri" Institute for Pharmacological Research,
Milan, Italy
SOURCE: Cytokine+ (2005), 30(3), 125-131
CODEN: CYTIE9; ISSN: 1043-4666
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Infiltration of polymorphonuclear neutrophils (PMNs) is thought to play a role in ischemic brain damage. The present study investigated the effect of repertaxin, a new noncompetitive allosteric inhibitor for the receptors of the inflammatory chemokine CXC ligand 8 (CXCL8)/interleukin-8 (IL-8), on PMN infiltration and tissue injury in rats. Cerebral ischemia was induced by permanent or transient occlusion of the middle cerebral artery and myeloperoxidase activity, a marker of PMN infiltration, and infarct volume were evaluated 24 h later. Repertaxin (15 mg/kg) was administered systemically at the time of ischemia and every 2 h for four times. In permanent ischemia repertaxin reduced PMN infiltration by 40% in the brain cortex but did not limit tissue damage. In transient ischemia (90-min ischemia followed by reperfusion), repertaxin inhibited PMN infiltration by 54% and gave 44% protection from tissue damage. Repertaxin had anti-inflammatory and neuroprotective effects also when given at reperfusion and even at 2 h of reperfusion. The protective effect of repertaxin did not interfere with brain levels of the chemokine. Since the PMN infiltration and its inhibition by repertaxin were comparable in the two models we conclude that reperfusion induces PMN activation, and inhibition of CXCL8 by repertaxin might be of pharmacol. interest in transient ischemia.

OS.CITING REF COUNT: 19 THERE ARE 19 CAPLUS RECORDS THAT CITE THIS RECORD (19 CITINGS)
REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 17 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:201863 CAPLUS
DOCUMENT NUMBER: 142:385080

TITLE: Predicting Human Serum Albumin Affinity of
Interleukin-8 (CXCL8) Inhibitors by 3D-QSPR Approach
Aureli, Loretta; Cruciani, Gabriele; Cesta, Maria
Candida; Anacardio, Roberto; De Simone, Lucio;
Moriconi, Alessio

CORPORATE SOURCE: Molecular Discovery Ltd., London, W1A 3BQ, UK
SOURCE: Journal of Medicinal Chemistry (2005), 48(7),
2469-2479
CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 142:385080

AB A novel class of 2-(R)-phenylpropionamides has been recently reported to inhibit in vitro and in vivo interleukin-8 (CXCL8)-induced biol. activities. These CXCL8 inhibitors are derivs. of phenylpropionic nonsteroidal antiinflammatory drugs (NSAIDs), high-affinity ligands for site II of human serum albumin (HSA). Up to date, only a limited number of in silico models for the prediction of albumin protein binding are available. A three-dimensional quant. structure-property relationship (3D-QSPR) approach was used to model the exptl. affinity constant (Ki) to plasma proteins of 37 structurally related molcs., using physicochem. and 3D-pharmacophoric descriptors. Mol. docking studies highlighted that training set molcs. preferentially bind site II of HSA. The obtained model shows satisfactory statistical parameters both in fitting and predicting validation. External validation confirmed the statistical significance of the chemometric model, which is a powerful tool for the prediction of HSA

binding in virtual libraries of structurally related compds.
OS.CITING REF COUNT: 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS
RECORD (14 CITINGS)
REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 18 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2005:28032 CAPLUS
DOCUMENT NUMBER: 142:190637
TITLE: Inhibition of interleukin-8 (CXCL8/IL-8) responses by
repertaxin, a new inhibitor of the chemokine receptors
CXCR1 and CXCR2
AUTHOR(S): Casilli, Federica; Bianchini, Andrea; Gloaguen,
Isabelle; Biordi, Leda; Alesse, Edoardo; Festuccia,
Claudio; Cavalieri, Barbara; Strippoli, Raffaele;
Cervellera, Maria Neve; Di Bitondo, Rosa; Ferretti,
Elisabetta; Mainiero, Fabrizio; Bizzarri, Cinzia;
Colotta, Francesco; Bertini, Riccardo
CORPORATE SOURCE: Dompe S.p.A. Research Center, L'Aquila, Italy
SOURCE: Biochemical Pharmacology (2005), 69(3), 385-394
CODEN: BCPCA6; ISSN: 0006-2952
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Repertaxin is a new non-competitive allosteric blocker of interleukin-8
(CXCL8/IL-8) receptors (CXCR1/R2), which by locking CXCR1/R2 in an
inactive conformation prevents receptor signaling and human
polymorphonuclear leukocyte (PMN) chemotaxis. Given the unique mode of
action of repertaxin it was important to examine the ability of repertaxin
to inhibit a wide range of biol. activities induced by CXCL8 in human
leukocytes. Our results show that repertaxin potently and selectively
blocked PMN adhesion to fibrinogen and CD11b up-regulation induced by
CXCL8. Reduction of CXCL8-mediated PMN adhesion by repertaxin was paralleled
by inhibition of PMN activation including secondary and tertiary granule
release and pro-inflammatory cytokine production, whereas PMN phagocytosis of
Escherichia coli bacteria was unaffected. Repertaxin also selectively
blocked CXCL8-induced T lymphocyte and natural killer (NK) cell migration.
These data suggest that repertaxin is a potent and specific inhibitor of a
wide range of CXCL8-mediated activities related to leukocyte recruitment
and functional activation in inflammatory sites.
OS.CITING REF COUNT: 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS
RECORD (22 CITINGS)
REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 19 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2004:803495 CAPLUS
DOCUMENT NUMBER: 141:343217
TITLE: Repertaxin, a novel inhibitor of rat CXCR2 function,
inhibits inflammatory responses that follow intestinal
ischaemia and reperfusion injury
AUTHOR(S): Souza, Danielle G.; Bertini, Riccardo; Vieira,
Angelica T.; Cunha, Fernando Q.; Poole, Steve;
Allegretti, Marcello; Colotta, Francesco; Teixeira,
Mauro M.
CORPORATE SOURCE: Immunopharmacology, Departamento de Bioquímica e
Imunologia, ICB, Universidade Federal de Minas Gerais,
Belo Horizonte, Brazil
SOURCE: British Journal of Pharmacology (2004), 143(1),
132-142
CODEN: BJPCBM; ISSN: 0007-1188
PUBLISHER: Nature Publishing Group

DOCUMENT TYPE: Journal
LANGUAGE: English

AB Neutrophils are thought to play a major role in the mediation of reperfusion injury. CXC chemokines are known inducers of neutrophil recruitment. Here, we assessed the effects of Repertaxin, a novel low mol. weight inhibitor of human CXCL8 receptor activation, on the local, remote and systemic injuries following intestinal ischemia and reperfusion (I/R) in the rat. Pre-incubation of rat neutrophils with Repertaxin (10⁻¹¹-10⁻⁶ M) inhibited the chemotaxis of neutrophils induced by human CXCL8 or rat CINC-1, but not that induced by fMLP, PAF or LTB₄, in a concentration-dependent manner. Repertaxin also prevented CXCL8-induced calcium

influx but not CXCL8 binding to purified rat neutrophils. In a model of mild I/R injury (30 min of ischemia and 30 min of reperfusion), Repertaxin dose-dependently (3-30 mg kg⁻¹) inhibited the increase in vascular permeability and neutrophil influx. Maximal inhibition occurred at 30 mg kg⁻¹. Following severe I/R injury (120 min of ischemia and 120 min of reperfusion), Repertaxin (30 mg kg⁻¹) markedly prevented neutrophil influx, the increase in vascular permeability both in the intestine and the lungs. Moreover, there was prevention of hemorrhage in the intestine of reperfused animals. Repertaxin effectively suppressed the increase in tissue (intestine and lungs) and serum concns. of TNF- α and the reperfusion-associated lethality. For comparison, we also evaluated the effects of an anti-CINC-1 antibody in the model of severe I/R injury. Overall, the antibody effectively prevented tissue injury, systemic inflammation and lethality. However, the effects of the antibody were in general of lower magnitude than those of Repertaxin. In conclusion, CINC-1 and possibly other CXC chemokines, acting on CXCR2, have an important role during I/R injury. Thus, drugs, such as Repertaxin, developed to block the function of the CXCR2 receptor may be effective at preventing reperfusion injury in relevant clin. situations.

OS.CITING REF COUNT: 38 THERE ARE 38 CAPLUS RECORDS THAT CITE THIS RECORD (38 CITINGS)
REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 20 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:703810 CAPLUS

DOCUMENT NUMBER: 141:343408

TITLE: Noncompetitive allosteric inhibitors of the inflammatory chemokine receptors CXCR1 and CXCR2: Prevention of reperfusion injury

AUTHOR(S): Bertini, Riccardo; Allegretti, Marcello; Bizzarri, Cinzia; Moriconi, Alessio; Locati, Massimo; Zampella, Giuseppe; Cervellera, Maria N.; di Cioccio, Vito; Cesta, Maria C.; Galliera, Emanuela; Martinez, Fernando O.; di Bitondo, Rosa; Troiani, Giulia; Sabbatini, Vilma; D'Anniballe, Gaetano; Anacardio, Roberto; Cutrin, Juan C.; Cavallieri, Barbara; Mainiero, Fabrizio; Strippoli, Raffaele; Villa, Pia; di Girolamo, Maria; Martin, Franck; Gentile, Marco; Santoni, Angela; Corda, Daniela; Poli, Giuseppe; Mantovani, Alberto; Ghezzi, Pietro; Colotta, Francesco
CORPORATE SOURCE: Dompe, L'Aquila, 67100, Italy

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (2004), 101(32), 11791-11796
CODEN: PNASA6; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The chemokine CXC ligand 8 (CXCL8)/IL-8 and related agonists recruit and activate polymorphonuclear cells by binding the CXC chemokine receptor 1

(CXCR1) and CXCR2. Here the authors characterize the unique mode of action of a small-mol. inhibitor (repertaxin) of CXCR1 and CXCR2. Structural and biochem. data are consistent with a noncompetitive allosteric mode of interaction between CXCR1 and repertaxin, which, by locking CXCR1 in an inactive conformation, prevents signaling. Repertaxin is an effective inhibitor of polymorphonuclear cell recruitment in vivo and protects organs against reperfusion injury. Targeting the repertaxin interaction site of CXCR1 represents a general strategy to modulate the activity of chemoattractant receptors.

OS.CITING REF COUNT: 75 THERE ARE 75 CAPLUS RECORDS THAT CITE THIS RECORD (75 CITINGS)
 REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 21 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:498365 CAPLUS

DOCUMENT NUMBER: 141:173953

TITLE: Acylmethanesulfonamides as new acylating agents for primary amines

AUTHOR(S): Coniglio, Silvia; Aramini, Andrea; Cesta, M. Candida; Colagiola, Sandro; Curti, Roberto; D'Alessandro, Fabrizio; D'Anniballe, Gaetano; D'Elia, Valerio; Nano, Giuseppe; Orlando, Valerie; Allegretti, Marcello

CORPORATE SOURCE: Dompe Research and Development, Chemistry Department, Dompe S.p.A., L'Aquila, 67100, Italy

SOURCE: Tetrahedron Letters (2004), 45(28), 5375-5378

CODEN: TELEAY; ISSN: 0040-4039

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:173953

AB A simple and efficient procedure for the preparation of secondary amides through internal condensation of acylmethanesulfonamides ammonium salts is described. The selective acylation of mixed primary-secondary amines could be an attractive application of this method.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 22 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:615394 CAPLUS

DOCUMENT NUMBER: 137:150277

TITLE: Use of (R)-ibuprofen methanesulfonamide and salts thereof in the treatment and prevention of ischemia/reperfusion injury or rejection reactions of transplanted organs

INVENTOR(S): Bertini, Riccardo; Colotta, Francesco; Novellini, Roberto

PATENT ASSIGNEE(S): Dompe S.p.A., Italy

SOURCE: PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002062330	A2	20020815	WO 2002-EP946	20020130
WO 2002062330	A3	20030403		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
 RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
 UZ, VN, YU, ZA, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,
 GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA,
 GN, GQ, GW, ML, MR, NE, SN, TD, TG

CA 2432432	A1	20020815	CA 2002-2432432	20020130
CA 2432432	C	20080325		
AU 2002250869	A1	20020819	AU 2002-250869	20020130
AU 2002250869	B2	20061019		
EE 200300340	A	20031015	EE 2003-340	20020130
EP 1355641	A2	20031029	EP 2002-719742	20020130
EP 1355641	B1	20050921		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
HU 2003003024	A2	20031229	HU 2003-3024	20020130
HU 2003003024	A3	20050728		
BR 2002006804	A	20040203	BR 2002-6804	20020130
JP 2004517948	T	20040617	JP 2002-562337	20020130
CN 1561205	A	20050105	CN 2002-804226	20020130
NZ 526655	A	20050225	NZ 2002-526655	20020130
RU 2257895	C2	20050810	RU 2003-126600	20020130
AT 304846	T	20051015	AT 2002-719742	20020130
ES 2248541	T3	20060316	ES 2002-719742	20020130
IL 157180	A	20090615	IL 2002-157180	20020130
ZA 2003004861	A	20040630	ZA 2003-4861	20030623
NO 2003003273	A	20030718	NO 2003-3273	20030718
KR 857898	B1	20080910	KR 2003-709714	20030723
MX 2003006686	A	20040531	MX 2003-6686	20030725
US 20040102520	A1	20040527	US 2003-250465	20031002
US 7560487	B2	20090714		

PRIORITY APPLN. INFO.: IT 2001-MI206 A 20010202
 WO 2002-EP946 W 20020130

AB The use of (R)-ibuprofen methanesulfonamide is described for the preparation of medicaments for the treatment and prevention of ischemia/reperfusion injury or functional injury resulting from rejection reactions of transplanted organs. In particular, the use of non-toxic salts of (R)-ibuprofen methanesulfonamide, such as the (L)-lysine salt (DF 1681B), is described for the prevention and the treatment of rejection reactions of transplanted kidneys. DF 1681B prevented renal function impairment secondary to cold ischemia in a rat model of kidney transplantation.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 23 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2000:290989 CAPLUS

DOCUMENT NUMBER: 132:321722

TITLE: Preparation of N-(2-arylpropionyl)sulfonamides as inhibitors of neutrophil chemotaxis and degranulation induced by interleukin 8.

INVENTOR(S): Bertini, Riccardo; Bizzarri, Cinzia; Sabbatini, Vilma; Porzio, Stefano; Caselli, Gianfranco; Allegretti, Marcello; Cesta, Maria Candida; Gandolfi, Carmelo A.; Mantovanini, Marco; Colotta, Francesco

PATENT ASSIGNEE(S): Dompe' S.P.A., Italy; et al.

SOURCE: PCT Int. Appl., 41 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000024710	A1	20000504	WO 1999-EP7740	19991014
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
IT 1303249	B1	20001106	IT 1998-MI2280	19981023
CA 2347752	A1	20000504	CA 1999-2347752	19991014
BR 9914741	A	20010703	BR 1999-14741	19991014
EP 1123276	A1	20010816	EP 1999-953824	19991014
EP 1123276	B1	20030108		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
TR 200101124	T2	20011022	TR 2001-1124	19991014
HU 2001003793	A2	20020328	HU 2001-3793	19991014
HU 2001003793	A3	20030929		
HU 225107	B1	20060628		
EE 200100233	A	20020815	EE 2001-233	19991014
EE 4912	B1	20071015		
JP 2002528434	T	20020903	JP 2000-578281	19991014
JP 4194761	B2	20081210		
AT 230723	T	20030115	AT 1999-953824	19991014
ES 2190264	T3	20030716	ES 1999-953824	19991014
NZ 511077	A	20030829	NZ 1999-511077	19991014
AU 769850	B2	20040205	AU 2000-10375	19991014
CN 1615833	A	20050518	CN 2004-10085635	19991014
RU 2255084	C2	20050627	RU 2001-113733	19991014
IL 142496	A	20060115	IL 1999-142496	19991014
CZ 296434	B6	20060315	CZ 2001-1441	19991014
CN 100368394	C	20080213	CN 1999-812451	19991014
SK 286372	B6	20080805	SK 2001-538	19991014
MX 2001003987	A	20020225	MX 2001-3987	20010420
NO 2001002000	A	20010620	NO 2001-2000	20010423
US 6887903	B1	20050503	US 2001-830075	20011121
HK 1041255	A1	20080718	HK 2002-102780	20020412
NZ 525084	A	20040827	NZ 2003-525084	20030401
US 20030216392	A1	20031120	US 2003-460203	20030613
US 6881755	B2	20050419		
AU 2003259648	A1	20031127	AU 2003-259648	20031103
AU 2003259648	B2	20060525		
EP 1579859	A1	20050928	EP 2004-7177	20040325
EP 1579859	B1	20061213		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
AT 347883	T	20070115	AT 2004-7177	20040325
ES 2279248	T3	20070816	ES 2004-7177	20040325
AU 2005226901	A1	20051006	AU 2005-226901	20050317
CA 2555162	A1	20051006	CA 2005-2555162	20050317
WO 2005092315	A1	20051006	WO 2005-EP2822	20050317
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM,
 SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
 MR, NE, SN, TD, TG

CN 1933825	A	20070321	CN 2005-80009560	20050317
BR 2005009167	A	20070911	BR 2005-9167	20050317
JP 2007530478	T	20071101	JP 2007-504310	20050317
MX 2006009085	A	20070402	MX 2006-9085	20060810
KR 2007018015	A	20070213	KR 2006-718031	20060905
NO 2006004793	A	20061023	NO 2006-4793	20061023
US 20090030083	A1	20090129	US 2006-588454	20061205
PRIORITY APPLN. INFO.:			IT 1998-MI2280	A 19981023
			AU 2000-10375	A3 19991014
			WO 1999-EP7740	W 19991014
			US 2001-830075	A3 20011121
			EP 2004-7177	A 20040325
			WO 2005-EP2822	W 20050317

OTHER SOURCE(S): MARPAT 132:321722

AB R2CHMeCONR1SO2R (R2 = aryl; R = alkyl, CF3, cyclohexyl, o-tolyl,
 3-pyridyl, 2-pyridylethyl, p-cyanophenylmethyl, p-aminophenylmethyl,
 3-cyano-1-Pr, 4-aminobutyl, etc.; R1 = H, alkyl), were prepared Thus,
 (R)-2-(4-isobutylphenyl)propionyl chloride in MeCN was added to NH3 in H2O
 at 0-5° to give (R)-2-(4-isobutylphenyl)propionamide. Title
 compds. inhibited chemotaxis of PMN human leukocytes with IC50 = 10-7 to
 10-9M.

OS.CITING REF COUNT: 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS
 RECORD (13 CITINGS)
 REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

---Logging off of STN---

Connection closed by remote host
 END

Unable to generate the STN prompt.
 Exiting the script...

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:esptacrs1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page for STN Seminar Schedule - N. America
 NEWS 2 JAN 12 Match STN Content and Features to Your Information
 Needs, Quickly and Conveniently
 NEWS 3 JAN 25 Annual Reload of MEDLINE database
 NEWS 4 FEB 16 STN Express Maintenance Release, Version 8.4.2, Is
 Now Available for Download
 NEWS 5 FEB 16 Derwent World Patents Index (DWPI) Revises Indexing
 of Author Abstracts
 NEWS 6 FEB 16 New FASTA Display Formats Added to USGENE and PCTGEN
 NEWS 7 FEB 16 INPADOCDB and INPAFAMDB Enriched with New Content
 and Features
 NEWS 8 FEB 16 INSPEC Adding Its Own IPC codes and Author's E-mail
 Addresses
 NEWS 9 APR 02 CAS Registry Number Crossover Limits Increased to
 500,000 in Key STN Databases
 NEWS 10 APR 02 PATDPAFULL: Application and priority number formats
 enhanced
 NEWS 11 APR 02 DWPI: New display format ALLSTR available
 NEWS 12 APR 02 New Thesaurus Added to Derwent Databases for Smooth
 Sailing through U.S. Patent Codes
 NEWS 13 APR 02 EMBASE Adds Unique Records from MEDLINE, Expanding
 Coverage back to 1948
 NEWS 14 APR 07 CA/Caplus CLASS Display Streamlined with Removal of
 Pre-IPC 8 Data Fields
 NEWS 15 APR 07 50,000 World Traditional Medicine (WTM) Patents Now
 Available in Caplus
 NEWS 16 APR 07 MEDLINE Coverage Is Extended Back to 1947
 NEWS EXPRESS FEBRUARY 15 10 CURRENT WINDOWS VERSION IS V8.4.2,
 AND CURRENT DISCOVER FILE IS DATED 15 JANUARY 2010.
 NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN customer
 agreement. This agreement limits use to scientific research. Use
 for software development or design, implementation of commercial
 gateways, or use of CAS and STN data in the building of commercial
 products is prohibited and may result in loss of user privileges
 and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 11:55:43 ON 12 MAY 2010

```
=> file registry
COST IN U.S. DOLLARS                SINCE FILE      TOTAL
                                   ENTRY      SESSION
FULL ESTIMATED COST                0.22          0.22
```

FILE 'REGISTRY' ENTERED AT 11:55:53 ON 12 MAY 2010
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2010 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
 provided by InfoChem.

STRUCTURE FILE UPDATES: 11 MAY 2010 HIGHEST RN 122257-16-0

DICTIONARY FILE UPDATES: 11 MAY 2010 HIGHEST RN 1222257-16-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

ISCA INFORMATION NOW CURRENT THROUGH January 8, 2010.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> file caplus medline	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	1.47	1.69

FILE 'CAPLUS' ENTERED AT 11:57:35 ON 12 MAY 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 11:57:35 ON 12 MAY 2010

=> s bertini r?/au or colotta f?/au
L1 536 BERTINI R?/AU OR COLOTTA F?/AU

=> s l1 and spinal cord
L2 2 L1 AND SPINAL CORD

=> dup rem l2
PROCESSING COMPLETED FOR L2
L3 1 DUP REM L2 (1 DUPLICATE REMOVED)

=> d l3 ibib abs

L3 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2010 ACS on STN DUPLICATE 1
ACCESSION NUMBER: 2007:976827 CAPLUS
DOCUMENT NUMBER: 147:314799
TITLE: Reparixin, an inhibitor of CXCR2 function, attenuates inflammatory responses and promotes recovery of function after traumatic lesion to the spinal cord
AUTHOR(S): Gorio, Alfredo; Madaschi, Laura; Zadra, Giorgia; Marfia, Giovanni; Cavallieri, Barbara; Bertini, Riccardo; Di Giulio, Anna Maria
CORPORATE SOURCE: Pharmacological Laboratories, Department of Medicine, Surgery and Dentistry, Faculty of Medicine, University of Milan, Milan, Italy
SOURCE: Journal of Pharmacology and Experimental Therapeutics (2007), 322(3), 973-981
CODEN: JPETAB; ISSN: 0022-3565
PUBLISHER: American Society for Pharmacology and Experimental Therapeutics
DOCUMENT TYPE: Journal
LANGUAGE: English
AB It has been shown that the blockade of CXCR1 and CXCR2 receptors prevents ischemia/reperfusion damage in several types of vascular beds. Reparixin

is a recently described inhibitor of human CXCR1/R2 and rat CXCR2 receptor activation. We applied reparixin in rats following traumatic spinal cord injury and determined therapeutic temporal and dosage windows. Treatment with reparixin significantly counteracts secondary degeneration by reducing oligodendrocyte apoptosis, migration to the injury site of neutrophils and ED-1-pos. cells. The observed preservation of the white matter might also be secondary to the enhanced proliferation of NG2-pos. cells. The expression of macrophage-inflammatory protein-2, tumor necrosis factor- α , interleukin (IL)-6, and IL-1 β was also counteracted, and the proliferation of glial fibrillary acidic protein-pos. cells was markedly reduced. These effects resulted in a smaller post-traumatic cavity and in a significantly improved recovery of hind limb function. The best beneficial outcome of reparixin treatment required 7-day administration either by i.p. route (15 mg/kg) or s.c. infusion via osmotic pumps (10 mg/kg), reaching a steady blood level of 8 μ g/mL. Methylprednisolone was used as a reference drug; such treatment reduced cytokine production but failed

to affect the rate of hind limb recovery.

OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD
(8 CITINGS)
REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s interleukin 8 or il-8 or cxcl8
L4 46473 INTERLEUKIN 8 OR IL-8 OR CXCL8

=> s l4 and spinal cord injury
L5 28 L4 AND SPINAL CORD INJURY

=> s l5 not py>2004
L6 11 L5 NOT PY>2004

=> dup rem l6
PROCESSING COMPLETED FOR L6
L7 10 DUP REM L6 (1 DUPLICATE REMOVED)

=> d l7 ibib abs 1-10

L7 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2010 ACS on STN
ACCESSION NUMBER: 2004:570498 CAPLUS
DOCUMENT NUMBER: 141:94383
TITLE: Composition and method to prevent and treat brain and
spinal cord injuries
INVENTOR(S): Wang, Yanming
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 8 pp., Cont.-in-part of U.S.
Ser. No. 962,009.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040138125	A1	20040715	US 2003-744114	20031223
US 20030059476	A1	20030327	US 2001-962009	20010924
US 6683066	B2	20040127		

PRIORITY APPLN. INFO.: US 2001-962009 A2 20010924
AB A composition and method for treating and preventing injury to central nervous

system tissue are provided. The composition is comprised of agents that can increase colloidal osmotic pressure and osmolality. The method comprises withdrawing cerebrospinal fluid from the subarachnoid spaces around the tissue to be treated or protected and injecting the composition into subarachnoid spaces. The acute spinal cord ischemia was induced in 28 rabbits. In the group of rabbits treated with the composition no apparent deficit could be observed, and all of rabbits walked and moved smoothly.

L7 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2010 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2004:84128 CAPLUS

DOCUMENT NUMBER: 140:246868

TITLE: Antithrombin reduces the ischemia/reperfusion-induced spinal cord injury in rats by attenuating inflammatory responses

AUTHOR(S): Hirose, Koji; Okajima, Kenji; Taoka, Yuji; Uchiba, Mitsuhiro; Nakano, Kan-yu; Utoh, Junichi; Kitamura, Nobuo

CORPORATE SOURCE: Departments of Surgery and Diagnostic Medicine, Graduate School of Medical Science, Kumamoto University, Kumamoto, Japan

SOURCE: Thrombosis and Haemostasis (2004), 91(1), 162-170
CODEN: THHAQD; ISSN: 0340-6245

PUBLISHER: Schattauer GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Antithrombin (AT) reveals its antiinflammatory activity by promoting endothelial release of prostacyclin (PGI₂) in vivo. Since neuroinflammation is critically involved in the development of ischemia/reperfusion (I/R)-induced spinal cord injury (SCI), it is possible that AT reduces the I/R-induced SCI by attenuating the inflammatory responses. We examined this possibility using rat model of I/R-induced SCI in the present study. AT significantly reduced the mortality and motor disturbances by inhibiting reduction of the number of motor neurons in animals subjected to SCI. Microinfarctions of the spinal cord seen after reperfusion were markedly reduced by AT. AT significantly enhanced the I/R-induced increases in spinal cord tissue levels of 6-keto-PGF_{1α}, a stable metabolite of PGI₂. AT significantly inhibited the I/R-induced increases in spinal cord tissue levels of TNF- α , rat interleukin-8 and myeloperoxidase. In contrast, Trp49-modified AT did not show any protective effects. Pretreatment with indomethacin significantly reversed the protective effects of AT. An inactive derivative of factor Xa, which selectively inhibits thrombin generation, has been shown to fail to reduce SCI. Taken together, these observations strongly suggested that AT might reduce I/R-induced SCI mainly by the antiinflammatory effect through promotion of endothelial production of PGI₂. These findings also suggested that AT might be a potential neuroprotective agent.

OS.CITING REF COUNT: 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:334388 CAPLUS

DOCUMENT NUMBER: 138:352740

TITLE: Immortalized human microglia cell and continuous cell line for screening therapeutics for treatment of autoimmune and neurodegenerative diseases

INVENTOR(S): Kim, Seung U.

PATENT ASSIGNEE(S): Can.

SOURCE: U.S. Pat. Appl. Publ., 28 pp., Cont.-in-part of U.S. Ser. No. 855,468.

DOCUMENT TYPE: CODEN: USXXCO
 LANGUAGE: Patent
 FAMILY ACC. NUM. COUNT: English
 PATENT INFORMATION: 3

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030082139	A1	20030501	US 2001-887145	20010622
US 6780641	B2	20040824		
WO 2002004618	A1	20020117	WO 2000-US18777	20000710
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 20020064877	A1	20020530	US 2001-855468	20010515
CA 2383942	A1	20020117	CA 2001-2383942	20010709
WO 2002004604	A2	20020117	WO 2001-IB1770	20010709
W: AU, CA, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				

PRIORITY APPLN. INFO.: WO 2000-US18777 A2 20000710
 US 2001-855468 A2 20010515
 US 2001-887145 A 20010622
 WO 2001-IB1770 W 20010709

AB An immortalized human cell line is provided which has the characteristics of human embryonic microglia. Such immortalized microglia cells express CD68, CD11c and MHC class I and II antigens as surface markers; have demonstrable phagocytic properties; and produce progeny continuously while maintained in culture. A method of transforming human microglial cells into an immortalized cell line is also provided. The genetically modified human microglia cells can express active substances from a selected group consisting of MIP-1 β , MCP-1, IL-1 β , IL-6, IL-12, and IL-15; and in the stimulated state can overexpress at least cytokines, chemokines, and other cytotoxic and neurotoxic substances. Such immortalized microglia cells can be used for screening of compds. for diseases. These cells may be utilized for the treatment of at least Alzheimer disease, Parkinson disease, Huntington disease, amyotrophic lateral sclerosis, stroke, spinal cord injuries, ataxia, autoimmune diseases and AIDS-dementia.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2010 ACS ON STN
 ACCESSION NUMBER: 2003:241782 CAPLUS
 DOCUMENT NUMBER: 138:260460
 TITLE: Pharmaceutical composition for brain and spinal cord injuries
 INVENTOR(S): Wang, Yanming
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 9 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

US 20030059476	A1	20030327	US 2001-962009	20010924
US 6683066	B2	20040127		
WO 2003026565	A2	20030403	WO 2002-US28918	20020911
WO 2003026565	A3	20031120		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002335736	A1	20030407	AU 2002-335736	20020911
US 20040142905	A1	20040722	US 2003-703320	20031107
US 20040142906	A1	20040722	US 2003-703830	20031107
US 20040138125	A1	20040715	US 2003-744114	20031223
PRIORITY APPLN. INFO.:			US 2001-962009	A 20010924
			WO 2002-US28918	W 20020911

AB After interruption of blood supply to central nervous system tissue, cerebral edema sets in. It has been shown that restoring blood flow to injured areas of the central nervous system after the onset of edema does not result in blood reperfusion of the tissue. A composition and method for treating injured central nervous tissue, or preventing injury to central nervous system tissue is provided. The composition is generally an amphipathic lipid in an oil solution. The method provides for withdrawing cerebrospinal fluid from the subarachnoid spaces around the tissue to be treated or protected, and replacing that fluid with an approx. equivalent volume of the amphipathic lipid in oil composition. The treatment can be augmented with agents that suppress production of cerebrospinal fluid, or with other known agents. Acute spinal cord ischemia was induced in 29 rabbits. At both 24 h and 1 wk after ischemia, in group 1 (soybean oil treatment) and group 2 [vitamin E injection solution (1 mg/mL)] treatment all rabbits showed no behavioral deficit, all walked and moved smoothly. In group 3 (12 rabbits for control), all of rabbits showed complete spastic paraplegia with no movement to the hind limbs.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L7 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2002:51606 CAPLUS
 DOCUMENT NUMBER: 136:113767
 TITLE: Immortalized human microglia cell and continuous cell line containing exogenous genes and uses in therapy
 INVENTOR(S): Kim, Seung U.
 PATENT ASSIGNEE(S): University of British Columbia, Can.
 SOURCE: PCT Int. Appl., 46 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
WO 2002004604	A2	20020117	WO 2001-IB1770	20010709
W: AU, CA, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
WO 2002004618	A1	20020117	WO 2000-US18777	20000710

W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE,
 DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
 KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN,
 MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
 TR, TT, UA, UG, US, UZ, VN, YU, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
 CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

US 20020064877	A1	20020530	US 2001-855468	20010515
US 20030082139	A1	20030501	US 2001-887145	20010622
US 6780641	B2	20040824		
CA 2383942	A1	20020117	CA 2001-2383942	20010709

PRIORITY APPLN. INFO.:

WO 2000-US18777	W	20000710
US 2001-855468	A	20010515
US 2001-887145	A	20010622
WO 2001-IB1770	W	20010709

AB An immortalized human cell line is provided which has the characteristics of human embryonic microglia. Such immortalized microglia cells express CD68, CD11c and MHC class I and II antigens as surface markers; have demonstrable phagocytic properties; and produce progeny continuously while maintained in culture. A method of transforming human microglial cells into an immortalized cell line is also provided. The genetically modified human microglia cells can express active substances from a selected group consisting of MIP-1 β , MCP-1, IL-1 β , IL-6, IL-12, and IL-15; and in the stimulated state can overexpress at least cytokines, chemokines, and other cytotoxic and neurotoxic substances. Such immortalized microglia cells can be used for screening of compds. for diseases. These cells may be utilized for the treatment of at least Alzheimer disease, Parkinson disease, Huntington disease, amyotrophic lateral sclerosis, stroke, spinal cord injuries, ataxia, autoimmune diseases and AIDS-dementia.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD
 (1 CITINGS)

L7 ANSWER 6 OF 10 MEDLINE on STN
 ACCESSION NUMBER: 2001505893 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 11553680
 TITLE: A neutrophil elastase inhibitor (ONO-5046) reduces
 neurologic damage after spinal cord
 injury in rats.
 AUTHOR: Tonai T; Shiba K; Taketani Y; Ohmoto Y; Murata K; Muraguchi
 M; Ohsaki H; Takeda E; Nishisho T
 CORPORATE SOURCE: Department of Orthopedic Surgery and Clinical Research
 Institute, National Zentsuji Hospital, Kagawa, Japan..
 ttonai@zentsuji.hosp.go.jp
 SOURCE: Journal of neurochemistry, (2001 Sep) Vol. 78, No. 5, pp.
 1064-72.
 Journal code: 2985190R. ISSN: 0022-3042. L-ISSN: 0022-3042.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200110
 ENTRY DATE: Entered STN: 17 Sep 2001
 Last Updated on STN: 29 Oct 2001
 Entered Medline: 25 Oct 2001

AB In view of a cytoprotective effect of elastase inhibitor on
 chemokine-mediated tissue injury, we examined the neuroprotective effect
 of ONO-5046, a specific inhibitor of neutrophil elastase, in rats with
 spinal cord injury. Standardized spinal cord
 compression markedly increased cytokine-induced neutrophil

chemo-attractant (CINC)-1 mRNA and protein. Their increases correlated with neurologic severity of injured rats. Immunohistochemically, CINC-1 protein was detected sequentially in vascular endothelial cells at 4 h, in perivascular neutrophils at 8 h, and in neutrophils infiltrating into cord substance at 12 h. Pretreatment with ONO-5046 (50 mg/kg) markedly ameliorated motor disturbance in injured rats, and reduced CINC-1 protein and mRNA expression. ONO-5046 also significantly reduced the increase of neutrophil accumulation or infiltration estimated by myeloperoxidase activity, and the extent of vascular permeability by Evans blue extravasation in the injured cord segment in comparison to control animals receiving vehicle. These results suggest that CINC-1 contributed to inflammation in rat spinal cord injury and ONO-5046 attenuated neurologic damage partly by blocking CINC-1 production of the chemoattractant, preventing neutrophil activation and vascular endothelial cell injury.

L7 ANSWER 7 OF 10 MEDLINE on STN
 ACCESSION NUMBER: 2001168114 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 11269455
 TITLE: Proinflammatory cytokines in cerebrospinal fluid in repair of thoracoabdominal aorta.
 AUTHOR: Kuniyoshi T; Sasaki S; Shiiya N; Miyatake T; Mafune N; Yasuda K
 CORPORATE SOURCE: Department of Cardiovascular Surgery, Hokkaido University School of Medicine, Sapporo, Japan..
 SOURCE: kuniyoshi@med.hokudai.ac.jp
 The Annals of thoracic surgery, (2001 Mar) Vol. 71, No. 3, pp. 801-6.
 Journal code: 15030100R. ISSN: 0003-4975. L-ISSN: 0003-4975.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: (COMPARATIVE STUDY)
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals
 ENTRY MONTH: 200105
 ENTRY DATE: Entered STN: 4 Jun 2001
 Last Updated on STN: 4 Jun 2001
 Entered Medline: 31 May 2001

AB BACKGROUND: Little is known about alterations of cytokine levels in cerebrospinal fluid (CSF) during thoracoabdominal aortic surgery. We measured perioperative CSF cytokine levels to determine their clinical significances. METHODS: Perioperative serum and CSF levels of cytokine were measured in 15 adult patients undergoing repair of the descending thoracic aorta (n = 4) or thoracoabdominal aorta (n = 11). All patients underwent prosthetic replacement and perioperative CSF drainage. Serum and CSF levels of tumor necrosis factor-alpha, Interleukin- (IL-) 1beta, IL-6, IL-8, IL-10, and IL-12 were measured before operation and at 0, 6, 12, 18, 24, 48, and 72 hours postoperatively using enzyme-linked immunosorbent assays. RESULTS: There were no hospital deaths, but 1 patient suffered paraplegia. Cerebrospinal fluid IL-8 levels peaked at immediately after operation (751.7 +/- 42.1 pg/mL versus preoperative levels, 54.9 +/- 24.6 pg/mL; p < 0.001), and the higher levels persisted for 72 hours. In contrast, serum IL-8 levels did not change and remained lower than CSF levels. The patient with paraplegia had the highest CSF IL-8 levels throughout the study period. Serum and CSF levels of tumor necrosis factor-alpha, IL-1beta, IL-6, and IL-12 did not significantly change. Serum and CSF levels of IL-10 were significantly elevated after operation compared with preoperative levels. In contrast to IL-8, serum IL-10 levels surpassed CSF levels. CONCLUSIONS: Cerebrospinal fluid IL-8 levels are significantly

elevated in thoracoabdominal aortic operation, and may be the most sensitive to the inflammatory response in the ischemic spinal cord injury. Persistent elevation of CSF IL-8 levels may be predictive of further development of neurologic deficits, and a reduction of proinflammatory cytokine levels may be a beneficial effect of CSF drainage, but this requires further investigation.

L7 ANSWER 8 OF 10 MEDLINE on STN
ACCESSION NUMBER: 2000212801 MEDLINE
DOCUMENT NUMBER: PubMed ID: 10750763
TITLE: Lazaroid reduces production of IL-8 and IL-1 receptor antagonist in ischemic spinal cord injury.
AUTHOR: Kunihara T; Sasaki S; Shiiya N; Ishikura H; Kawarada Y; Matsukawa A; Yasuda K
CORPORATE SOURCE: Department of Cardiovascular Surgery, Hokkaido University School of Medicine, Sapporo, Japan.
SOURCE: The Annals of thoracic surgery, (2000 Mar) Vol. 69, No. 3, pp. 792-8.
Journal code: 15030100R. ISSN: 0003-4975. L-ISSN: 0003-4975.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, NON-U.S. GOV'T)
LANGUAGE: English
FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals
ENTRY MONTH: 200004
ENTRY DATE: Entered STN: 5 May 2000
Last Updated on STN: 5 May 2000
Entered Medline: 27 Apr 2000

AB BACKGROUND: 21-aminosteroids (Lazaroids) have demonstrated the protective effect against cerebral ischemic injury through the inhibition of lipid peroxidation. We examined whether Lazaroids affected the production of proinflammatory and antiinflammatory cytokines in ischemic spinal cord injury model. MATERIALS: Anesthetized New Zealand white rabbits underwent a 20-minute infrarenal aortic cross-clamping (AXC) with pretreatment of either an intravenous 3 mg/kg Lazaroid U74389G (group L; n = 10) or the same volume saline (group P; n = 10). Sham operation group (group S; n = 6) underwent only exposure of the aorta. Plasma concentrations of interleukin (IL)-8, -1beta, -1 receptor antagonist (IL-1ra) and tumor necrosis factor (TNF)-alpha were measured at four time points. Functional assessment with Tarlov score at 24 and 48 hours after pretreatment, pathologic assessment of the spinal cord, and measurements of cytokine levels in the spinal cord were performed. RESULTS: The maximum elevation of plasma IL-8 and -1ra levels occurred at 1 hour after declamping in four measurement points. Plasma IL-8 and -1ra levels in group L were significantly lower than those in group P (*p < 0.05). Plasma TNFalpha peaked at 5 minutes after declamping, but decreased afterwards. Plasma TNFalpha levels were not different among three groups. Spinal IL-8 levels in group L (0.98 +/- 0.34 ng/g tissue) were lower than those in group P (7.26 +/- 2.26 ng/g tissue) (*p < 0.05). Spinal IL-1ra and TNFalpha were not significantly different. Tarlov score and pathologic assessment were better in group L. CONCLUSIONS: Lazaroid U-74389G reduced the production of systemic IL-8 and -1ra and spinal IL-8 when AXC caused spinal cord injury. These results indicate that Lazaroids may attenuate ischemic endothelial cell injury or activation of leukocytes.

L7 ANSWER 9 OF 10 MEDLINE on STN

ACCESSION NUMBER: 2000408569 MEDLINE
DOCUMENT NUMBER: PubMed ID: 10903607
TITLE: Activated protein C reduces the
ischemia/reperfusion-induced spinal cord
injury in rats by inhibiting neutrophil activation.
AUTHOR: Hirose K; Okajima K; Taoka Y; Uchiba M; Tagami H; Nakano K;
Utoh J; Okabe H; Kitamura N
CORPORATE SOURCE: First Department of Surgery and the Department of
Laboratory Medicine, Kumamoto University School of
Medicine, Kumamoto, Japan.
SOURCE: Annals of surgery, (2000 Aug) Vol. 232, No. 2, pp. 272-80.
Journal code: 0372354. ISSN: 0003-4932. L-ISSN: 0003-4932.
Report No.: NLM-PMC1421140.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals
ENTRY MONTH: 200008
ENTRY DATE: Entered STN: 1 Sep 2000
Last Updated on STN: 1 Sep 2000
Entered Medline: 22 Aug 2000

AB OBJECTIVE: To examine whether activated protein C (APC) reduces
spinal cord injury in rats by inhibiting
neutrophil activation after the transient ischemia. SUMMARY BACKGROUND
DATA: Ischemic spinal cord injury is an
important pathologic mechanism leading to the paraplegia observed after
surgery to repair aortic aneurysms. Activated neutrophils play a pivotal
role in the development of ischemia/reperfusion-induced tissue injury.
Recently, the authors have reported that APC, a physiologic anticoagulant,
prevents lipopolysaccharide-induced pulmonary vascular injury by
inhibiting neutrophil activation. These observations strongly suggest
that APC reduces ischemia/reperfusion-induced spinal
cord injury by inhibiting neutrophil activation.
METHODS: In rats, spinal cord ischemia was induced by using a balloon
catheter placed into the aorta. After the transient ischemia, survival
and motor function were evaluated, and histologic examination of the
spinal cord was performed by using both hematoxylin-and-eosin staining and
2,3,5, -triphenyltetrazolium chloride (TTC) staining 24 hours after the
ischemia. Tissue levels of myeloperoxidase and cytokines, including tumor
necrosis factor-alpha (TNF-alpha) and rat interleukin-8
, were measured in six experimental groups: sham-operated, control, APC
(100 microg/kg, intravenous), dansyl glutamyl-glycyl-arginyl chloromethyl
ketone-treated activated factor X (DEGR-F.Xa), a selective inhibitor of
thrombin generation (1 mg/kg, intravenous), nitrogen mustard-induced
leukocytopenia, and diisopropyl fluorophosphate-treated APC (DIP-APC),
active site-blocked APC (100 microg/kg, intravenous). APC, DEGR-F.Xa, and
DIP-APC were administered intravenously 30 minutes before aortic
occlusion. Control and leukocytopenic rats received saline instead of
other drugs. RESULTS: Pretreatment with APC significantly reduced motor
disturbances compared with those in control animals. In contrast, neither
DEGR-F.Xa nor DIP-APC had any effect. Microinfarctions, evidenced by the
absence of TTC staining and histologic change, were markedly reduced in
animals given APC. The increases in the tissue levels of TNF-alpha, rat
interleukin-8, and myeloperoxidase in the ischemic part
of the spinal cord were significantly reduced in animals that received
APC. These levels were not reduced in rats given DEGR-F.Xa or DIP-APC.
Leukocytopenia produced effects similar to those of APC. CONCLUSIONS: APC
reduced the ischemia/reperfusion-induced spinal cord
injury by inhibiting neutrophil activation. The therapeutic
mechanisms of APC might depend on its inhibitory effect on the production
of TNF-alpha, which is a potent activator of neutrophils. Although the
anticoagulant effects of APC might not be related to its ability to

inhibit TNF-alpha production, its serine protease activity appears to be essential in the therapeutic mechanism. APC appears to have potential as a therapeutic agent for prevention of spinal cord injury in patients undergoing aortic aneurysm repair.

L7 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 1998:527446 CAPLUS
 DOCUMENT NUMBER: 129:145631
 ORIGINAL REFERENCE NO.: 129:29595a,29598a
 TITLE: Expression vectors with ubiquitin promoter and methods for in vivo expression of therapeutic polypeptides
 INVENTOR(S): Johansen, Teit E.
 PATENT ASSIGNEE(S): Neurosearch A/S, Den.; Bavarian Nordic Research Institute A/S
 SOURCE: PCT Int. Appl., 29 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9832869	A1	19980730	WO 1998-DK37	19980129
W: JP, US				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP 961830	A1	19991208	EP 1998-900847	19980129
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				

PRIORITY APPLN. INFO.: DK 1997-102 A 19970129
 WO 1998-DK37 W 19980129

AB The present invention relates to recombinant expression vectors carrying a gene encoding a therapeutically active polypeptide, which gene is under transcriptional control of a ubiquitin promoter. The invention also relates to the use of a ubiquitin promoter to direct in vivo expression of therapeutic genes after transfer of such genes to the central nervous system. The expression vectors include herpes virus vectors, vaccinia virus vectors, adeno-associated virus vectors, retroviral vectors, and adenovirus vectors. Vector-expressed therapeutic genes may encode a nerve growth factor, a fibroblast growth factor, an insulin-like growth factor, etc.

OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
56.46	58.15

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-5.95	-5.95

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 12:02:22 ON 12 MAY 2010